



FUNGICIDE

# Cover XL®

Cover XL® is a broad-spectrum, preventative fungicide with systemic and curative properties recommended for the control of many important plant diseases. Cover XL® may improve the yield and/or quality of the crop due to the positive effects on plant physiology. Cover XL® may be applied as a foliar spray in alternating spray programs or in tank mixes with other crop protection products.

## PRODUCT FEATURES:

- Cover XL® will travel throughout the plant by means of translocation to provide preventative and curative benefits.
- Applications of Cover XL® can be made by ground, air or through chemigation providing long-lasting residual control of a variety of diseases.
- Cover XL® may be applied for early-season disease control and beneficial physiological effects in certain crops.

## CROP TOLERANCE

Plant tolerance has been found to be acceptable for all crops on the label, however, not all possible tank mix combinations have been tested under all conditions. When possible, it is recommended to test the combinations on a small portion of the crop to ensure that a phytotoxic response will not occur because of application. See Product Use Precautions on the label for apple phytotoxicity information.

ACTIVE INGREDIENT: Azoxystrobin and Propiconazole

FUNGICIDE GROUP: 11 & 3

ACTIVE INGREDIENT PER GALLON: 1.18 lbs Azoxystrobin and 1.02 lbs Propiconazole

MODE OF ACTION: QoI Inhibitor and Demethylation Inhibitor

USE RATES: 7 – 28 ounces

PACKAGING: 2 x 2.5's and 265 gal



**Albaugh®**  
Your Alternative®

ALBAUGHLLC.COM



## CORN (Field, and Pop (Includes Seed Production))

TARGET DISEASE	Use Rate fl. oz. product/A	COMMENTS
Anthracnose Leaf Blight ( <i>Colletotrichum graminicola</i> ) Eye Spot ( <i>Aureobasidium zeae</i> ) Gray Leaf Spot ( <i>Cercospora zeae-maydis</i> ) Northern Corn Leaf Blight ( <i>Setosphaeria turcica</i> ) Northern Corn Leaf Spot ( <i>Cochliobolus carbonum</i> ) Physoderma Brown Spot ( <i>Physoderma maydis</i> ) Rusts ( <i>Puccinia</i> spp.) Southern Corn Leaf Blight ( <i>Cochliobolus heterostrophus</i> ) also known as Helminthosporium Leaf Blights ( <i>H. maydis</i> , <i>H. turcicum</i> , <i>H. carbonum</i> ) Suppression of: Diplodia Ear Rot ( <i>D. maydis</i> )	10.5	Early application (V4-V8): An early application (V4-V8) of Cover XL® may be applied for early season disease control and plant performance benefits. If mixing with herbicides other than solo glyphosate products, consult your local Albaugh representative.
	10.5 - 14	Later season applications: For gray leaf spot, rusts, anthracnose, and eye spot, apply 10.5-14 oz./A Cover XL® when disease first appears. If conditions favorable for disease persist, continue to apply on a 14-day schedule.  For leaf blights apply 10.5-14 oz. Cover XL® when disease first appears. Continue on a 7- to 14-day schedule. Use the low rate when disease pressure is low. Under heavy disease pressure or if conditions are favorable for disease, apply the high rate.  Do not use adjuvants or other additives after the V8 growth stage and prior to the VT growth stage, as use during these development times may impose stress on the plant that could inhibit proper kernel development. VT is defined as when the last branch of the tassel is completely visible, but silks have not yet emerged from the ear shoot.  Apply no more than 2 applications of Cover XL® or any other Group 11 fungicide per year. Use of an adjuvant such as COC may provide additional disease control.
APPLICATION: For best results, sufficient coverage is very important. For ULV aerial applications DO NOT use less than 1.0 GPA. Use a higher water volume for aerial application if equipment and/or conditions will not provide good coverage. Cover XL® may be applied by ground, air (ULV), or chemigation.		
FIELD and POP CORN USE RESTRICTIONS:	Do not apply more than 56 fl. oz./A/season of Cover XL®. Do not apply more than 28 fl. oz. (0.224 lb. a.i. propiconazole) for field corn harvested for forage. Do not apply more than 0.45 lb. a.i. propiconazole-containing products/A/season. Do not apply within 30 days of harvest (30-day PHI) for forage, grain, or stover.	

## SOYBEANS (Soybean, Immature Seed (edamame))

TARGET DISEASE	Use Rate fl. oz. product/A	COMMENTS
Aerial Blight ( <i>Rhizoctonia solani</i> ) Alternaria Leaf Spot ( <i>Alternaria</i> spp.) Anthracnose ( <i>Colletotrichum truncatum</i> ) Brown Spot ( <i>Septoria glycines</i> ) Cercospora Blight and Leaf Spot ( <i>Cercospora kikuchii</i> ) Frogeye Leaf Spot ( <i>Cercospora soja</i> ) Pod and Stem Blight ( <i>Diaporthe phaseolorum</i> ) Soybean Rust ( <i>Phakopsora pachyrhizi</i> )	10.5 - 21	Foliar diseases (except rust): Apply 14-21 fl. oz./A at growth stage R3 (early pod set) when pods are inch long and 14-21 days later at growth stage R5 (pod fill). Cover XL® may be applied earlier should conditions be conducive for disease.  Soybean Rust: Apply 14-21 fl. oz./A at first indication that disease is in the area. For best control, preventive applications work best. Repeat on a 14- to 21-day interval. Use higher rate and shorter interval when diseases are present in the field and incidence is less than 2% (2 plants in 100 are infected). If incidence is greater than this or if disease is in mid-canopy, control will not be acceptable. Scouting for the disease and/or being aware of the proximity of the disease via monitoring systems will aid in the proper timing to maximize the effectiveness of the fungicide applications.  On certain varieties, Cover XL® applications may cause crinkled, smaller and/or greener leaves. Yields of beans displaying these characteristics have not been reduced due to Cover XL® treatments.
APPLICATION: Cover XL® is most effective when applied and allowed to dry before a rainfall. For best results, sufficient coverage is very important. DO NOT use less than 2.0 GPA. Use a higher water volume for aerial application if equipment and/or conditions will not provide for good coverage. Cover XL® may be applied by ground, air or chemigation.		
RESTRICTIONS	Do not apply more than 42 fl. oz./A of Cover XL® per crop. Do not apply more than 0.34 lb. a.i. of propiconazole-containing products/A/season. Do not apply more than 1.5 lb. a.i. of azoxystrobin-containing products/A/season. Apply up to Stage R6.	

Always read and follow label instructions  
 EPA Reg. No. 42750-289 | AD062215