

Contains Trinexapac-ethyl, the active ingredient used in Palisade® EC.



## PLANT GROWTH REGULATOR

# PINZOLA<sup>TM</sup> EC



## DECREASE LODGING, INCREASE STABILITY

Adverse weather and high nitrogen levels can contribute to detrimental lodging, posing a serious threat to cereal grower's investments. Pinzola<sup>TM</sup> EC uses the active ingredient trinexapac-ethyl to shorten nodes of cereal plants through inhibition of cell elongation. As a result, lodging potential is decreased and stability of plants is increased. See results in barley, wheat, sugarcane, rye, rice and more. Plus, the added benefit of a wide application window and tank-mix compatibility allows growers flexibility with efficient spray programs.

### KEY BENEFITS

- Increases stem thickness and diameter to help strengthen the stem and decrease lodging
- Wide application window gives growers flexibility with efficient spray programs
- Improves crop stability by decreasing crop height

### KEY USES

- Barley
- Grasses (grown for seed)
- Oats
- Rice
- Rye
- Triticale
- Wheat

### PRODUCT NOTES

#### EPA REGISTRATION NUMBER

91234-179

#### ACTIVE INGREDIENT

Trinexapac-ethyl 12%

#### FORMULATION

Emulsifiable Concentrate

#### HRAC NUMBER

23

#### SIGNAL WORD

Warning

#### PACKAGE SIZE

2 x 2.5 gal

#### RESTRICTED USE

No

Refer to label for specific use restrictions.



Relevant. Simple. Reliable.



## APPLICATION INFORMATION

Pinzola EC may be applied with all types of spray equipment commonly used for making ground and aerial applications. Proper adjustments and calibration of spraying equipment to give good coverage is essential for good growth regulator effects.

### Ground Application:

Apply in a minimum of 10 gallons of water per acre, unless specified otherwise. Do not apply through any ultra-low volume (ULV) spray system. Thorough coverage is necessary for good growth regulator effects.

### Aerial Application:

Thorough coverage is necessary to provide a good, uniform effect. A minimum of 2 gallons of diluent per acre can be used in grasses grown for seed, cereals, and sugarcane. Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Do not apply through any ultra-low volume (ULV) spray system.

### Application Through Irrigation Systems (Chemigation):

Apply this product only through center pivot, solid set, hand move, or moving wheel irrigation systems. Do not apply this product through any other type of irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. Apply in 0.1 - 0.25 inches/A of water. Excessive water may reduce efficacy. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

*Refer to product label for complete application and mixing instructions.*

©2019 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state. Pinzola™ EC is not manufactured, or distributed by Syngenta, seller of Palisade® EC. Palisade® EC is a registered trademark of Syngenta Group Company. Pinzola™ is a trademark of Atticus, LLC.



[www.AtticusLLC.com](http://www.AtticusLLC.com)