



HERBICIDE

# Priority™ 8E

Priority™ 8E is a selective herbicide for use as a preplant surface-applied, preplant incorporated or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in corn (field, pop, sweet and grown for seed), cotton, peanuts, soybeans, sorghum and Crop Group 6 legume vegetable (succulent and dried), potatoes, safflowers and tomatoes. Priority™ 8E can also be applied as a postemergent treatment on corn, cotton, potatoes, soybeans and tomatoes.

## PRODUCT FEATURES

- Priority™ 8E is an excellent choice as part of an integrated weed control program by providing broad spectrum weed control in a variety of crops.
- Priority™ 8E when mixed with other EPA registered herbicides can increase the spectrum of weed control helping to minimize weed resistance.
- Priority™ 8E can be applied through ground, aerial or chemigation application equipment.

ACTIVE INGREDIENT: Priority™

HERBICIDE GROUP: 15

ACTIVE INGREDIENT PER GALLON: 7.64 lbs

MODE OF ACTION: Long-chain Fatty Acid Inhibitor


USE RATES: See label for all product use rates

PACKAGING: 2 x 2.5's and 265 gals



Albaugh®  
Your Alternative®

ALBAUGHLLC.COM



## SOYBEANS – Priority™ 8E ALONE

Apply Priority™ 8E, either preplant surface-applied, preplant incorporated, preemergence, or post emergence using the appropriate rate specified below.

**Preplant Surface-Applied, Preplant Incorporated, Preemergence, or Postemergence:** Follow instructions for use of this product alone under Application Procedures on the product label.

### PREPLANT SURFACE-APPLIED FALL APPLICATION

- Apply after September 30 in MN, ND, SD, WI, and North of Route 30 in IA
- Apply after October 15 North of Route 91 in NE and South of Route 30 in IA
- Apply after October 31 North of Route 136 in IL

In all locations, apply Priority™ 8E to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 to 2 pts./A on medium-textured and 2 pts./A on fine-textured soils. Do not apply to frozen ground. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not exceed an incorporation depth greater than 2 to 3 inches.

Minimize furrow and ridge formation in the tillage operations.

### Preplant Surface – Spring Application

Use on medium and fine soils with minimum-tillage or no-tillage systems in CO, CT, DE, IA, IL, IN, KS, KY, MA, MD, ME, MI, MN, MO, MT, ND, NE, NH, NY, OH, PA, RI, SD, TN, VA, VT, WI, WV, and WY. Apply two-thirds of the specified rate of Priority™ 8E (1.67 pts./A on medium soils and 2 pts./A on fine soils) as a split treatment 30 to 45 days prior to planting and the remainder at planting. Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts./A on coarse soils not more than 2 weeks before planting.

**Preplant Incorporated or Preemergence:** On coarse soils, apply 1 to 1.33 pts./A of Priority™ 8E if organic matter content is less than 3%, or 1.33 pts./A if organic matter content is 3% or greater. On medium soils, apply 1.33 to 1.67 pts./A of this product. On fine soils, apply 1.33 to 1.67 pts./A of this product if organic matter content is less than 3%, or 1.67 to 2 pts./A if organic matter content is 3% or greater.

On soybeans, Priority™ 8E may be used up to 2.75 pts./A as a preplant surface-applied, preplant incorporated, or preemergence treatment on soils having an organic matter

content between 6% and 20%. The total rate of this product applied to soybeans during any one crop season must not exceed 2.75 pts./A.

### RESTRICTIONS: (PREPLANT SURFACE APPLIED, PREPLANT INCORPORATED, PREEMERGENCE):

- The total Priority™ 8E rate applied to must not exceed 2.75 pints / acre per year.
- If a pre-plant surface, pre-plant incorporated or pre-emergence application of metolachlor products has already been applied, a postemergence application of this product cannot be used.
- DO NOT graze or feed treated soybean forage, hay, or straw to livestock 30 days following treatment, or illegal residues may result.
- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate on soybeans, or illegal residues may result.
- DO NOT apply to frozen ground.

### POSTEMERGENCE APPLICATIONS

**Postemergence (From emergence up through the 3rd trifoliate leaf stage)**

For postemergence treatments, apply 1.0 to 1.33 pints per acre to soybeans from emergence through the 3rd trifoliate leaf stage. This product will not control emerged weeds, it must be applied to a weed free surface or in tank mixture with products that provide postemergence control of weeds present at the time of application.

Make postemergence applications at least 90 days before harvest.

### RESTRICTIONS (POSTEMERGENCE)

- To prevent illegal residues, DO NOT apply more than 1.33 pints per acre postemergence.
- DO NOT graze or feed treated soybean forage or soybean hay to livestock following a postemergence application of this product;
- DO NOT apply a postemergence application of this product if a pre-plant surface, pre-plant incorporated or pre-emergence application of metolachlor products has already been applied
- NOT FOR USE IN CALIFORNIA.

## SOYBEANS – Priority™ 8E COMBINATIONS

### Preplant Incorporated or Pre-emergence

Refer to product label for all restrictions and rate limitations for Priority™ 8E and tank mix partner products.

Always read and follow label directions.  
EPA Reg No. 42750-340 | AD050820  
Priority™ is a trademark of Albaugh, LLC