FUNGICIDE



PROVEN, LONG-LASTING RESIDUAL CONTROL

Slant™ is a broad-spectrum fungicide that provides a full-season of control from crop damaging diseases, helping growers achieve maximum yield potential. Slant delivers both curative and preventative activity, protecting crops at every stage.


Slant moves systemically, protecting new growth with long-lasting residual control. It works like a growth inhibitor, putting a stop to the spread of harmful diseases. Slant is an excellent tank-mix partner, offering maximum flexibility and efficiency.

KEY BENEFITS

- Protective, curative and systemic activity
- Full-season control
- · Protects new growth
- Excellent tank-mix partner
- · Low use rates

KEY USES

- Almonds
- Pecans
- Barley
- Rice
- Corn
- $\bullet \; Sorghum$
- Oats
- Soybeans
- Peanuts
- Wheat



PRODUCT NOTES

EPA REGISTRATION NUMBER 91234-25

ACTIVE INGREDIENT

Propiconazole 41.8%

FORMULATION

Emulsifiable Concentrate

FRAC NUMBER

3

SIGNAL WORD

Caution

PACKAGE SIZE

2 x 2.5 gal

135 gal

265 gal

RESTRICTED USE

No



Refer to product label for complete list.





APPLICATION INFORMATION

Slant is most effective when applied and allowed to dry before a rainfall. Avoid applying Slant under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply in a manner which results in exposure to humans or animals.

GROUND APPLICATION

For tree crops, apply Slant in a minimum of 50 gallons of water per acre unless otherwise specified. For all other crops, apply Slant in a minimum of 10 gallons of water per acre unless otherwise specified.

ARIAL APPLICATION

For tree crops, apply Slant in a minimum of 10 gallons of water per acre unless otherwise specified. For all other crops, apply Slant in a minimum of 2 gallons of water per acre unless otherwise specified.

CHEMIGATION APPLICATION

Apply Slant through properly equipped chemigation systems for disease control in the labeled crops. Refer to crop specific use directions for application rates, timing and frequency of application. Do not apply Slant by chemigation to other labeled crops except as specified in the product labeling. When applying this product by chemigation, do not exceed labeled rates or apply more frequently than recommended for conventional application methods. Slant, alone or in combination with other pesticides that are registered for application through irrigation systems, may be applied through irrigation systems. For chemigation application to labeled crops, apply in 0.1 to 0.25 inches of water unless otherwise specified. Chemigation with excessive water may lead to a decrease in efficacy.

MIXING INSTRUCTIONS

When preparing spray mixture, only prepare amount required for immediate use. For best results, thoroughly clean spray equipment before using this product. Agitate the spray solution before and during application.

APPLICATION RATE OF SLANT 41.8% EC (FL OZ/A)	ACTIVE INGREDIENT EQUIVALENT (LB A.I./A)	ACRES/GAL OF SLANT 41.8% EC
2.0	0.056	64.0
4.0	0.1125	32.0
6.0	0.169	21.3
8.0	0.225	16.0
10.0	0.28	12.8
12.0	0.34	10.7
16.0	0.45	8.0
20.0	0.56	6.4
24.0	0.67	5.3
30.0	0.84	4.3
32.0	0.90	4.0

Refer to product label for complete application and mixing instructions.

©2020 Atticus. Important: Always read and follow label instructions. Not all products are registered for use in every state.

Slant™ is not manufactured, or distributed by Syngenta, seller of Tilt® and Orbit®. Tilt® and Orbit® are registered trademarks of Syngenta Group Company. Slant™ is a trademark of Atticus, LLC.



KEY DISEASES

Aggregate sheath spot

Anthracnose

Black sheath spot Brown leaf spot Brown patch Dollar spot Early leaf spot **Ergot** Eye spot Foot rot Glume blotch Gray leaf spot Helminthosporium leaf blight Kernel smut Late leaf spot Leaf blight Leaf smut Narrow brown leaf spot Net blotch Northern corn leaf blight Northern corn leaf spot Powdery mildew Rust Sheath blight Sheath spot Southern stem rot Spot blotch Tan spot Web blotch

(Refer to product label for complete list)